

**TWO NEW SPECIES OF *OXYAGRION* SELYS, 1876,
WITH A DESCRIPTION OF FIVE NEW LARVAE
(ZYGOPTERA: COENAGRIONIDAE)**

J.M. COSTA¹, L.O. IRINEU DE SOUZA² and T.C. SANTOS¹

¹ Departamento, de Entomologia, Museu Nacional, Universidade Federal do Rio de Janeiro,
Quinta da Boa Vista, BR-20942 Rio de Janeiro, RJ, Brasil

² Universidade Federal de Mato Grosso do Sul, MS, Brasil

Received November 15, 1998 / Revised and Accepted June 12, 1999

O. pseudocardinale sp. n. (holotype ♂: Brazil, Minas Gerais, Fazenda da Cachoeira F.F. de Souza, 13-II-1990) and *O. sulmatogrossense* sp. n. (holotype ♂: Brazil, Mato Grosso do Sul, Campo Grande, Campus UFMS, 24-XII-1997) are described and illustrated. The larvae of *O. basale* Selys, 1876; *O. haematinum* Selys, 1876; *O. pavidum* Selys, 1876; *O. santosi* Martins, 1967 and *O. sulinum* Costa, 1978 are described and illustrated for the first time. Keys are provided for the known *Oxyagrion* spp. and for the known larvae.

INTRODUCTION

Nineteen species of *Oxyagrion* are known from the Neotropical region (COSTA, 1978, 1988; SANTOS, 1961) and 5 species are known from Mato Grosso and Mato Grosso do Sul States: *O. chapadense* Costa, 1978; *O. evanescens* Calvert, 1909; *O. impunctatum* Calvert, 1909; *O. microstigma* Selys, 1876 and *O. pavidum* Selys, 1876. Of the 19 species, 15 occur in Brazil and of these only *O. sulinum* Costa, 1978 had not been recorded from Mato Grosso State.

The two new species we describe here belong to *Oxyagrion* because they have red color patterns; lack postocular spots; black area on the head and on mesoepisternum is absent; penis with double fold; females with mesoepisternal fossae (except *O. hempelii* and *O. microstigma*) and with genital spine on 8th segment (except *O. evanescens*). All material, including types are lodged in the Museu Nacional, Rio de Janeiro.