

val, near the second striæ: rudimental striæ at the base, short and straight; epipleuræ with a longitudinal elevation near the base. Inhabits Georgia.

28. *CALATHIUS distinguendus*. Nob. Length $5\frac{1}{4}$ lines; breadth 2 lines.

Piceous black, shining, smooth; palpi, antennæ and legs ferruginous; front with a small impression at the base of the antennæ, and a linear one near the eyes: thorax smooth, with some indistinct transverse undulating wrinkles, quadrangular, narrowed before, the anterior angles projecting and a little rounded, base cut square, so as to make the posterior angles almost right, lateral margin rufo-piceous, a little curved, dilated, and depressed posteriorly; anteriorly; anterior transverse impression tolerably distinct: posterior well marked; medial line slight, abbreviated by the transverse impressions; with the anterior of which it forms an angle, so as to leave a well defined triangular depression in front; basal impressions slight, scarcely distinct from the depressed base of the thorax.

Elytra striate, striæ impunctured, with three impressed points on the third interval, the two upper ones near the third, the lower one close to the extremity of the second striæ. Inhabits Georgia.

Larger than *C. gregarius*, which it very much resembles; the thorax, and its margins are proportionally much wider, more curved, and more reflexed; and it wants the impressed point on each side of the thorax near the margin, which is very apparent in the *gregarius*: the anterior transverse impression is strongly marked, whereas in *C. gregarius*, it is much less apparent. It is also much darkened beneath.

29. *ANCHOMENUS Le Contei*. Dej. Cat. Length $9\frac{1}{2}$ lines; breadth 2 lines.

Very much resembles the *A. extensicollis*; but differs in the thorax being wider, and more rounded: the basal impressions are longer, perpendicular to the base, and curving upwardly and outwardly; whilst in the *extensicollis* they are oblique, so as to make them wider at the base; from this last circumstance the middle of the base between the impressions is much wider in *A. Le Contei* than in *A. extensicollis*.

It may, however, prove to be nothing more than a variety.

The Committee, to whom was referred the following paper, by Mr. Haldeman, reported in favour of publication:

Descriptions of Insects, presumed to be undescribed.

BY S. S. HALDEMAN.

LEUCOSPIS integra. Black, covered with pale yellowish pile; tegulæ and legs yellow. ♂ 5 millim. long.

♂ Antennæ yellowish beneath: thorax scabrous, with a spot of yellow pile posteriorly, wings translucent: legs yellow, except the basal half of the first and second pairs of femora, posterior femora black tip with yellow.

HEDYCHRUM janus. Brilliant, above violet, beneath uniform green; wings tip with fuliginous. 5 mill. long.